

DDR is located in the 97th MDG, RM 1262





DDR has an open door policy. Please stop by or call any time.

June 2016

Marijuana is the most commonly used illicit drug.

Delta-9-Tetrahydrocannabinol (THC) is the primary, active ingredient in cannabis psychoactive substances.

Marijuana and Hemp are cannabis derivatives.

Health Effects of Marijuana Use

- Respiratory illnesses
- Cancer Marijuana smoke contains 50-70% more carcinogens than tobacco smoke
- Short term memory damage
- Increased heart rate
- Impaired Coordination
- Increased anxiety
- Increased depression
- Increased suicidal ideation
- Difficulty problem solving
- Learning problems


Most Marijuana Use Begins in Adolescence: 78% of the 2.4 million people who began using in the last year were aged 12 to 20. 1

Of an estimated 113 million emergency room visits in the U.S. in 2006, marijuana was involved in 290,563 of these visits

A 2002 report concluded that the younger children are when they first use marijuana, the more likely they are to use cocaine and heroin and become chemically dependent.

Hemp-Based Nutritional

Supplements (Nutiva, Hempola, Manitoba After 5 days from last use, up to 20% of the THC from the dosage remains stored in body fat



Harvest etc.)

Per AFI 90-507, Alcohol and Drug Abuse Prevention and Treatment Program, the ingestion of hemp seed oil, hemp seed or products made with either is prohibited.

Failure to comply is a violation of Article 92, UCMJ.

It can take up to 30 days to eliminate a single dose of marijuana from an individual's body

While several states have legalized certain uses for Marijuana, the Department of Defense (DoD) maintains a zero tolerance policy and prohibits all marijuana use.

Chronic marijuana use, especially in a very young person, may also be a marker of risk for mental illnesses, including addiction, stemming from genetic or environmental vulnerabilities, such as early exposure to stress or violence.

High doses of marijuana can produce an acute psychotic reaction; in addition, use of the drug may trigger the onset or relapse of schizophrenia in vulnerable individuals.

^{1.} http://www.samhsa.gov/data/sites/default/files/NSDUHresultsPDFWHTML2013/Web/NSDUHresults2013.pdf